

Table 1 Larvicidal toxicity of seed methanolic extracts of *Asparagus setaceus* and *Cuscuta chinensis* on *Cx. quinquefasciatus*

Plant	Concentration (ppm)	Instars	Mean Mortality % \pm SE		
			24 h	48 h	72 h
<i>A.setaceus</i>	100	First	49.32 \pm 0.33	52.00 \pm 0.00	56.00 \pm 0.57
		Second	49.32 \pm 0.33	52.00 \pm 0.57	53.32 \pm 0.33
		Third	42.68 \pm 0.66	46.68 \pm 0.33	50.68 \pm 0.33
		Fourth	42.68 \pm 0.33	44.00 \pm 0.57	45.32 \pm 0.33
	control		00.00 \pm 0.00	00.00 \pm 0.00	00.00 \pm 0.00
	200	First	50.68 \pm 0.33	54.68 \pm 0.33	62.68 \pm 0.33
		Second	49.32 \pm 0.33	52.00 \pm 0.00	54.68 \pm 0.33
		Third	46.68 \pm 0.33	46.68 \pm 0.33	52.00 \pm 0.33
		Fourth	45.32 \pm 0.33	48.00 \pm 0.00	49.32 \pm 0.33
	control		00.00 \pm 0.00	00.00 \pm 0.00	00.00 \pm 0.00
	300	First	58.64 \pm 0.33	64.00 \pm 0.57	69.32 \pm 0.33
		Second	54.68 \pm 0.33	60.00 \pm 0.57	65.32 \pm 0.66
		Third	50.68 \pm 0.66	54.68 \pm 0.33	56.00 \pm 0.57
		Fourth	50.68 \pm 0.33	53.32 \pm 0.33	54.68 \pm 0.66
	control		00.00 \pm 0.00	00.00 \pm 0.00	00.00 \pm 0.00
	<i>C.chinensis</i>	100	First	54.68 \pm 0.33	57.32 \pm 0.33
Second			54.68 \pm 0.33	54.68 \pm 0.33	57.32 \pm 0.33
Third			50.68 \pm 0.33	52.00 \pm 0.57	57.32 \pm 0.66
Fourth			45.32 \pm 0.33	49.32 \pm 0.66	52.00 \pm 0.57
control			00.00 \pm 0.00	00.00 \pm 0.00	00.00 \pm 0.00
200		First	65.32 \pm 0.33	68.00 \pm 0.57	70.68 \pm 0.33
		Second	61.32 \pm 0.33	64.00 \pm 0.00	64.00 \pm 0.00
		Third	57.32 \pm 0.33	60.00 \pm 0.00	62.68 \pm 0.33
		Fourth	50.68 \pm 0.33	54.68 \pm 0.33	61.32 \pm 0.33
control			00.00 \pm 0.00	00.00 \pm 0.00	00.00 \pm 0.00
300		First	70.68 \pm 0.33	76.00 \pm 0.57	86.68 \pm 0.33
		Second	69.32 \pm 0.33	69.32 \pm 0.33	82.68 \pm 0.33
		Third	60.00 \pm 0.57	62.68 \pm 0.88	68.00 \pm 1.00
		Fourth	61.32 \pm 0.33	65.32 \pm 0.88	66.68 \pm 1.20
control			00.00 \pm 0.00	00.00 \pm 0.00	00.00 \pm 0.00