

Table 2 Weight loss and damage caused by *R. dominica* infestation on wheat seeds stored inside *E. aromatica*-treated jute bags at different doses after 42 days of storage

Doses(ml)	After 42 days			
	Number of adult emerged	%IR	% weight loss	% damage severity
2	3.20±0.32 ^b	90.34±1.23 ^b	1.62±0.08 ^b	5.11±0.67 ^b
4	1.44±0.88 ^a	92.00±0.63 ^b	0.84±0.03 ^a	3.54±0.21 ^a
6	0.00±0.00 ^a	100.00±0.00 ^b	0.00±0.00 ^a	0.00±0.00 ^a
8	0.00±0.00 ^a	100.00±0.00 ^b	0.00±0.00 ^a	0.00±0.00 ^a
Deep-soaked	0.00±0.00 ^a	100.00±0.00 ^b	0.00±0.00 ^a	0.00±0.00 ^a
Solvent control	18.24±0.34 ^c	0.00±0.00 ^a	34.86±0.67 ^c	22.33±0.28 ^c
Untreated control	19.65±1.05 ^c	0.00±0.00 ^a	34.92±0.45 ^c	23.89±0.88 ^c

Note: Each value is the mean ± standard error of seven replicates. Mean followed by the same letters within the same column are not significantly ($P > 0.05$) different from each other using New Duncan's Multiple Range Test