

Table 2 Time in days, the *An. arabiensis* Patton larvae took to develop to pupae and adult mosquitoes emerging from larval stages. Means are followed by SEM

Treatment	N	Crushed silver cyprinid fish				Tetramin [®] baby fish food				Control			
		Adult longevity	n	Time to pupation	n	Adult longevity	n	Time to pupation	n	Adult longevity	n	Time to pupation	n
Rain water alone	50	12.80±0.78	45	5.86±0.11 ^a	50	12.29±0.97	42	5.39±0.07 ^a	46	0	0	0	0
Doxycycline	50	9.00±1.00	2	10.00±2.00 ^a	3	10.00±1.64	18	7.32±0.15 ^a	28	0	0	0	0
Pollen	50	13.63±0.71	46	5.74±0.12 ^a	47	14.22±0.92	45	5.34±0.08 ^a	47	9.48±0.56	33	7.93±0.25	3
													3
Pollen + Doxycycline	50	10.31±1.43	13	8.30±0.34 ^a	20	9.86±0.84	29	6.78±0.19 ^a	36	0	0	0	0

Notes: 1 N is the total number of first instars larvae (L1s) used.

2 n is the number of pupae and adult mosquitoes per treatment,

3 Rows having mean pupation time superscripted with the same letters indicate a significant influence of the time it took the mosquito larval stage to pupate on adult mosquito longevity.