

Table 2 Total number (%) of mosquitoes collected on human bait during different months

Species	March	April	May	June	July	Total
<i>Cx. quinquefasciatus</i> *	55(13.68%)	54(13.43%)	117(29.1%)	114(28.36%)	62(15.42%)	402
<i>Cx. tritaeniorhynchus</i> #	10(19.61%)	11(21.51%)	16(31.37%)	8(15.69%)	6(11.76%)	51
<i>Cx. vishnui</i> #	-	-	5(45.45%)	3(27.27%)	3(27.27%)	11
<i>Cx. hutchinsoni</i>	8(57.14%)	6(42.86%)	-	-	-	14
<i>Ar. subalbatus</i> *	7(21.21%)	7(21.21%)	9(27.27%)	6(18.18%)	4(12.12%)	33
<i>Mn. uniformis</i> *	24(21.43%)	17(15.18%)	31(27.68%)	21(18.75%)	19(16.96%)	112
<i>Mn. annulifera</i> *	16(18.18%)	17(19.32%)	20(22.73%)	20(22.73%)	15(17.05%)	88
Total mosquitoes	120(16.88%)	112(15.75%)	198(27.85%)	172(24.19%)	109(15.33%)	711

Note: *Filariasis vectors, #Japanese encephalitis (JE) vectors