

Table 1 Total number (%) of mosquitoes collected on human bait during various phases of night (18h00 – 06h00)

Species	18h00-21h00	21h00-24h00	24h00-03h00	03h00-06h00	Total
<i>Cx. quinquefasciatus</i> *	67 (16.67%)	128 (31.84%)	149 (37.06%)	58 (14.43%)	402
<i>Cx. tritaeniorhynchus</i> #	6 (11.76%)	20 (39.23%)	19 (37.25%)	6 (11.76%)	51
<i>Cx. vishnui</i> #	1 (9.09%)	4 (36.36%)	5 (45.45%)	1 (9.09%)	11
<i>Cx. hutchinsoni</i>	1 (7.14%)	6 (42.86%)	5 (35.71%)	2 (14.29%)	14
<i>Ar. subalbatus</i> *	6 (18.18%)	8 (24.24%)	13 (39.39%)	6 (18.18%)	33
<i>Mn. uniformis</i> *	21 (18.75%)	37 (33.03%)	38 (33.93%)	16 (14.29%)	112
<i>Mn. annulifera</i> *	15 (17.04%)	27 (30.68%)	32 (36.36%)	14 (15.91%)	88
Total mosquitoes	117 (16.46%)	230 (32.35%)	261 (36.71%)	103 (14.49%)	711

Note: \*Filariasis vectors, #Japanese encephalitis (JE) vectors