

Table 2 Percent mortality of 3<sup>rd</sup> instars larvae of *Cx. vishnui* group using all the six solvents at 72 hours of post-exposure

Solvent extracts	Percent mortality ( Mean $\pm$ SE) at 72 Hours				
	20 ppm	40 ppm	60 ppm	80 ppm	100 ppm
Petroleum ether	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00	1.00 $\pm$ 0.00	1.00 $\pm$ 0.00
n-hexane	0.00 $\pm$ 0.00	1.2 $\pm$ 0.00	0.00 $\pm$ 0.00	0.85 $\pm$ 0.00	0.00 $\pm$ 0.00
Ethyl acetate	4.00 $\pm$ 0.72	4.00 $\pm$ 1.28	4.00 $\pm$ 0.75	4.00 $\pm$ 0.72	6.05 $\pm$ 0.25
Chloroform: methanol (1:1 v/v)	6.12 $\pm$ 0.28	6.12 $\pm$ 0.44	7.45 $\pm$ 0.88	7.45 $\pm$ 1.28	7.45 $\pm$ 0.45
acetone	35.00 $\pm$ 0.94	47.32 $\pm$ 0.82	68.00 $\pm$ 0.75	78.00 $\pm$ 0.00	95.32 $\pm$ 0.82
Absolute alcohol	0.00 $\pm$ 0.00	10.00 $\pm$ 0.94	16.68 $\pm$ 0.94	18.32 $\pm$ 0.94	22.00 $\pm$ 0.82