

Table 4: Source of Blood Meals in *Anopheles* mosquitoes collected in Mwea and Ahero study sites

Village	Anopheles species	Number of mosquitoes tested	Human Indoor	Human Outdoor	Cattle Indoor	Cattle Outdoor	Goat Indoor	Goat Outdoor	Dog Indoor	Dog Outdoor	Cat Indoor	Cat Outdoor	Chicken Indoor	Chicken Outdoor
Mbuinjeru	<i>An. arabiensis</i>	150	0.53	0.00	0.00	0.02	0.00	0.38	0.00	0.02	0.05	0.00	0.00	0.00
Ndindiruku	<i>An. arabiensis</i>	135	0.14	0.00	0.00	0.5	0.00	0.14	0.00	0.00	0.00	0.14	0.08	0.00
Murinduko	<i>An. arabiensis</i>	120	0.00	0.00	0.00	0.78	0.00	0.22	0.00	0.00	0.00	0.00	0.00	0.00
Kamagaga	<i>An. arabiensis</i>	20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	0.00
	<i>An. funestus</i>	110	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	0.00
Kobura	<i>An. arabiensis</i>	140	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	<i>An. funestus</i>	9	0.25	0.00	0.56	0.00	0.00	0.00	0.00	0.06	0.00	0.00	0.13	0.00
Wagai	<i>An. arabiensis</i>	9	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	0.00
	<i>An. funestus</i>	150	0.25	0.00	0.25	0.00	0.00	0.00	0.00	0.00	0.5	0.00	1.00	0.00