

Table 1 Effect of crude extract of mature fruit of *T. crispa* against 1st to 4th instars larvae of *Cx. quinquefasciatus*

Instar	Concentration (%)	Percentage mortality (Mean $\pm$ SE)		
		24 h	48 h	72 h
1 st	0.1	42.68 $\pm$ 0.67	62.68 $\pm$ 0.33	80.00 $\pm$ 0.58
	0.2	50.68 $\pm$ 0.33	69.32 $\pm$ 0.33	85.32 $\pm$ 0.33
	0.3	60.00 $\pm$ 0.58	80.00 $\pm$ 0.58	93.32 $\pm$ 0.33
	0.4	68.00 $\pm$ 0.58	89.32 $\pm$ 0.33	100.00 $\pm$ 0.00
	0.5	77.33 $\pm$ 0.33	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00
2 nd	0.1	38.68 $\pm$ 0.33	54.68 $\pm$ 0.33	72.00 $\pm$ 0.58
	0.2	45.32 $\pm$ 0.33	62.68 $\pm$ 0.33	78.68 $\pm$ 0.88
	0.3	50.67 $\pm$ 0.33	69.32 $\pm$ 0.33	88.00 $\pm$ 0.58
	0.4	58.68 $\pm$ 0.33	77.32 $\pm$ 0.33	96.00 $\pm$ 0.58
	0.5	66.68 $\pm$ 0.33	85.32 $\pm$ 0.33	100.00 $\pm$ 0.58
3 rd	0.1	30.68 $\pm$ 0.33	49.32 $\pm$ 0.33	68.00 $\pm$ 0.58
	0.2	38.68 $\pm$ 0.33	56.00 $\pm$ 0.58	76.00 $\pm$ 0.58
	0.3	44.00 $\pm$ 0.58	62.68 $\pm$ 0.33	81.32 $\pm$ 0.33
	0.4	52.00 $\pm$ 0.58	72.00 $\pm$ 0.58	89.32 $\pm$ 0.33
	0.5	60.00 $\pm$ 0.58	81.32 $\pm$ 0.33	100.00 $\pm$ 0.00
4 th	0.1	26.68 $\pm$ 0.33	42.68 $\pm$ 0.33	58.68 $\pm$ 0.33
	0.2	30.68 $\pm$ 0.33	46.68 $\pm$ 0.33	65.32 $\pm$ 0.33
	0.3	37.32 $\pm$ 0.33	53.32 $\pm$ 0.33	70.67 $\pm$ 0.33
	0.4	42.68 $\pm$ 0.33	58.68 $\pm$ 0.33	76.00 $\pm$ 0.58
	0.5	50.68 $\pm$ 0.33	66.67 $\pm$ 0.33	84.00 $\pm$ 0.58

Note: SE= Standard error