

Table 3 Number and percent of different species of man-landing (indoor and outdoor together) mosquitoes collected during August 2009 to July 2010 in tribal and non-tribal areas of Bankura district, West Bengal, India

Species	Tribal						Non-tribal					
	Day-biting		Night-biting		Total		Day-biting		Night-biting		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1. <i>Ar. subalbatus</i>	32	4.71	5	0.09	37	0.62	2617	69.29	560	3.67	3177	16.71
2. <i>An. annularis</i>	0	0.00	0	0.00	0	0.00	0	0.00	8	0.05	8	0.04
3. <i>An. barbirostris</i>	0	0.00	161	3.03	161	2.69	0	0.00	125	0.82	125	0.66
4. <i>An. subpictus</i>	39	5.74	163	3.07	202	3.37	20	0.53	139	0.91	159	0.84
5. <i>An. vagus</i>	179	26.32	1200	22.60	1379	23.03	97	2.57	1168	7.66	1265	6.65
6. <i>Cx. quinquefasciatus</i>	209	30.74	2144	40.38	2353	39.29	635	16.81	11739	77.02	12374	65.06
7. <i>Cx. vishnui</i> group	221	32.50	1628	30.66	1849	30.87	97	2.57	1278	8.39	1375	7.23
8. <i>Ma. indiana</i>	0	0.00	8	0.15	8	0.13	0	0.00	147	0.96	147	0.77
9. <i>St. aegypti</i>	0	0.00	0	0.00	0	0.00	244	6.46	63	0.41	307	1.61
10. <i>St. albopictus</i>	0	0.00	0	0.00	0	0.00	67	1.77	14	0.09	81	0.43
Total	680	100.00	5309	100.00	5989	100.00	3777	100.00	15241	100.00	19018	100.00